



1001 West Memorial, Suite 112  
 Oklahoma City, OK 73114  
 PH: (405) 509-6599  
 FlexCareInfusion.com

Please fax completed referral form to Flexcare:

**(888) 219-8102**

Your Infusion Specialist Field Contact: Cam Jones,

**(443) 221-9233**

# REMICADE INFUSION THERAPY REFERRAL FORM

PATIENT DEMOGRAPHICS:			
PATIENT NAME:	PREFERRED CONTACT #:		
DATE OF REFERRAL:	SECONDARY CONTACT #:		
SOCIAL SECURITY NUMBER:	ADDRESS:		
DATE OF BIRTH:	CITY, STATE, ZIP:		
PRIMARY DIAGNOSIS:			
ICD-IO CODE: <input type="checkbox"/> Crohn's Disease ICD- 10: _____	<input type="checkbox"/> Psoriatic Arthritis ICD- 10: _____		
<input type="checkbox"/> Ulcerative Colitis ICD- 10: _____	<input type="checkbox"/> Ankylosing Spondylitis ICD- 10: _____		
<input type="checkbox"/> Rheumatoid Arthritis ICD- 10: _____	<input type="checkbox"/> Plaque Psoriasis ICD- 10: _____		
PATIENT INFORMATION:			
ALLERGIES: <input type="checkbox"/> NKDA	FIRST DOSE: <input type="checkbox"/> Y <input type="checkbox"/> N		
	DATE OF LAST INFUSION:		
	NEXT DOSE DUE BY:		
HEIGHT: ____ Ft ____ In      WEIGHT: ____ Lb or ____ Kg	ACCESS/LINE TYPE: <input type="checkbox"/> PIV <input type="checkbox"/> PORT <input type="checkbox"/> PICC <input type="checkbox"/> MIDLINE		
GENDER: <input type="checkbox"/> F <input type="checkbox"/> M	OTHER:		
REQUIRED DOCUMENTATION: Please provide a copy of the following documents.			
<input checked="" type="checkbox"/> 1. INSURANCE CARD (Front & Back)	<input checked="" type="checkbox"/> 3. MOST RECENT LABS	<input checked="" type="checkbox"/> 5. NEGATIVE TB TEST RESULTS	
<input checked="" type="checkbox"/> 2. PATIENT DEMOGRAPHICS	<input checked="" type="checkbox"/> 4. H & P		
PRIMARY MEDICATION ORDER:		PRN & PREMEDICATIONS:	
<input type="checkbox"/> Remicade 3mg/kg IV at weeks 0, 2 and 6 and then every ____ weeks.	<b>MEDICATIONS</b>	<b>30 minutes prior every infusion</b>	<b>PRN</b>
<input type="checkbox"/> Remicade 5mg/kg IV at weeks 0, 2 and 6 and then every ____ weeks.	Acetaminophen ____ mg PO	<input type="checkbox"/>	<input type="checkbox"/> PRN every ____ hours for mild or moderate infusion reaction.
<input type="checkbox"/> Other orders (write other or additional orders below):	Diphenhydramine ____ mg PO	<input type="checkbox"/>	<input type="checkbox"/> PRN every ____ hours for mild or moderate infusion reaction.
	Diphenhydramine ____ mg diluted in 10mL 0.9% NaCl slow IV push over 2-3 minutes.	<input type="checkbox"/>	<input type="checkbox"/> PRN every ____ hours for mild or moderate infusion reaction.
If weight-based dosing, dose on the patients:	Methylprednisolone ____ mg IV push over 5 minutes.	<input type="checkbox"/>	
<input type="checkbox"/> ACTUAL BODY WEIGHT <input type="checkbox"/> IDEAL BODY WEIGHT	0.9% NaCl ____ mL to infuse over ____ minutes.	<input type="checkbox"/>	
Round up to nearest viral size?			
<input type="checkbox"/> YES <input type="checkbox"/> NO			
LINE USE/CARE ORDERS:		ADVERSE REACTION & ANAPHYLAXIS ORDERS:	
<input type="checkbox"/> START PIV/ACCESS CVC	<input type="checkbox"/> FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (See Reverse Side)	<input type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side)	
<input type="checkbox"/> OTHER FLUSH ORDERS:	<input type="checkbox"/> OTHER: (please fax other reaction orders if checking this box)		
PRESCRIBER INFORMATION:			
PHYSICIAN NAME:	PHONE:		
OFFICE CONTACT:	FAX:		
ADDRESS:	LICENSE #:		
CITY, STATE, ZIP:	NPI:		
▶▶▶▶▶▶▶▶▶▶▶▶▶▶▶▶	DATE:		
PHYSICIAN SIGNATURE:			

**FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:**

*\*This graph does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	<b>MILD INFUSION REACTION</b>	<b>MODERATE INFUSION REACTION</b>	<b>SEVERE INFUSION REACTION/ANAPHYLAXIS</b>
<b>SYMPTOM CLASSIFICATION</b>	<ul style="list-style-type: none"> <li>• Flushing</li> <li>• Dizziness</li> <li>• Headache</li> <li>• Apprehension</li> <li>• Diaphoresis</li> <li>• Palpitations</li> <li>• Nausea / Vomiting</li> <li>• Pruritis</li> </ul>	<ul style="list-style-type: none"> <li>• Chest Tightness</li> <li>• Shortness of Breath</li> <li>• Hypo/hypertension (&gt;20 mmHg Change in Systolic BP from Baseline)</li> <li>• Increased Temperature (&gt;2 Degrees Fahrenheit)</li> <li>• Urticaria</li> </ul>	<ul style="list-style-type: none"> <li>• Hypo/hypertension (&gt;40 mmHg Change in Systolic BP from Baseline).</li> <li>• Increase Temperature (&gt;2 Degrees Fahrenheit) with Rigors</li> <li>• Shortness of Breath with Wheezing</li> <li>• Laryngeal Edema</li> <li>• Chest Pain</li> <li>• Hypoxemia</li> </ul>
<b>TREATMENT PROTOCOL FOR ADULTS &gt;66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the mid- anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9%NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
<b>TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25 mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid- anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

**FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

**FLUSHING PROTOCOLS**

PATIENT CLASSIFICATION	LINE TYPE	FLUSHING PROTOCOL Normal Saline*			LOCKING PROTOCOL Heparin Sodium	
		PRE-ADMIN	POST ADMIN	POST LAB DRAW	10 Units/mL	100 Units/mL
<b>ADULT &gt;66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL	
	Midline	3 mL	3 mL		3 mL	
	Implanted Port	5 mL	10 mL	10 mL		5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL	
	Tunneled & non-Tunneled Catheters.	5 mL	10 mL	10 mL	5 mL	
<b>PEDIATRIC 33 LBS - 66LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL	
	Midline	3 mL	3 mL		3 mL	
	Implanted Port	5 mL	5 mL	10 mL	3 mL	
	Peripherally Inserted Central Catheter (PICC)	5 mL	5 mL	10 mL	3 mL	
	Tunneled & non-Tunneled Catheters.	5 mL	5 mL	10 mL	3 mL	

**FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

*\*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.*