

Please mark closest FlexCare location:

- Oklahoma City, OK     Norman, OK  
 Tulsa, OK             Birmingham, AL

**CONTACT INFORMATION**

Fax: (888) 219-8102

Email: orders@flexcareinfusion.com

Visit: flexcareinfusion.com/referrals

# OSTEOPOROSIS INFUSION THERAPY REFERRAL FORM

<b>PATIENT DEMOGRAPHICS:</b>	
PATIENT NAME:	PREFERRED CONTACT #:
DATE OF REFERRAL:	SECONDARY CONTACT #:
SOCIAL SECURITY NUMBER:	ADDRESS:
DATE OF BIRTH:	CITY, STATE, ZIP:

**PRIMARY DIAGNOSIS:**  
ICD-10 CODE:

<b>PATIENT INFORMATION:</b>	
ALLERGIES:    NKDA <input type="checkbox"/>	FIRST DOSE: <input type="checkbox"/> Y <input type="checkbox"/> N      NEXT DOSE DUE BY: _____
HEIGHT: ____ Ft ____ In      WEIGHT: ____ Lb or ____ Kg	DATE OF LAST INJECTION: _____
GENDER: <input type="checkbox"/> F <input type="checkbox"/> M	ACCESS/LINE TYPE: <input type="checkbox"/> PIV <input type="checkbox"/> PORT <input type="checkbox"/> PICC <input type="checkbox"/> MIDLINE

**REQUIRED DOCUMENTATION:**

Insurance Card (Front & Back)       Current Medication List       Bone Scan       Most Recent Labs Including Ca+ level within the past 3 months  
 Patient Demographics               H & P

PRIMARY MEDICATION ORDER:	PRN AND PREMEDICATIONS:		
<input type="checkbox"/> <b>PROLIA</b> 60mg subcutaneously, every 6 months <input type="checkbox"/> <b>EVENITY</b> 210mg subcutaneously monthly (2 x 105mg pre-filled syringes)	Acetaminophen ____ mg PO Diphenhydramine ____ mg PO Other: _____ _____ _____	Prior to every injection <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<b>PRN</b> <input type="checkbox"/> PRN every ____ hours after injection. <input type="checkbox"/> PRN every ____ hours after injection. <input type="checkbox"/> PRN every ____ hours after injection.

<b>LABS:</b> <input type="checkbox"/> Calcium <input type="checkbox"/> Other:  FREQUENCY: _____	<b>ADVERSE REACTION &amp; ANAPHYLAXIS ORDERS:</b> <input type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) *While reactions from injectable medications are rare, should a serious reaction occur, a clinician will obtain IV access and follow the referenced protocol)  <input type="checkbox"/> OTHER: (please fax other reaction orders if checking this box)
---	---

<b>PRESCRIBER INFORMATION:</b>	
PHYSICIAN NAME:	PHONE:
OFFICE CONTACT:	FAX:
ADDRESS:	LICENSE #:
CITY, STATE, ZIP:	NPI:
▶▶▶▶▶▶▶▶▶▶▶▶▶▶▶ PHYSICIAN SIGNATURE:	DATE:

# OSTEOPOROSIS INFUSION THERAPY REFERRAL FORM

## FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

*\*This graph does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
<b>SYMPTOM CLASSIFICATION</b>	<ul style="list-style-type: none"> <li>Flushing</li> <li>Dizziness</li> <li>Headache</li> <li>Apprehension</li> <li>Diaphoresis</li> <li>Palpitations</li> <li>Nausea / Vomiting</li> <li>Pruritis</li> </ul>	<ul style="list-style-type: none"> <li>Chest Tightness</li> <li>Shortness of Breath</li> <li>Hypo/hypertension (&gt;20 mmHg Change in Systolic BP from Baseline)</li> <li>Increased Temperature (&gt;2 Degrees Fahrenheit)</li> <li>Urticaria</li> </ul>	<ul style="list-style-type: none"> <li>Hypo/hypertension (&gt;40 mmHg Change in Systolic BP from Baseline).</li> <li>Increase Temperature (&gt;2 Degrees Fahrenheit) with Rigors</li> <li>Shortness of Breath with Wheezing</li> <li>Laryngeal Edema</li> <li>Chest Pain</li> <li>Hypoxemia</li> </ul>
<b>TREATMENT PROTOCOL FOR ADULTS &gt;66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the mid- anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9%NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
<b>TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25 mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid- anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

**FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

### FLUSHING PROTOCOLS

PATIENT CLASSIFICATION	LINE TYPE	FLUSHING PROTOCOL Normal Saline*			LOCKING PROTOCOL Heparin Sodium	
		PRE-ADMIN	POST ADMIN	POST LAB DRAW	10 Units/mL	100 Units/mL
<b>ADULT &gt;66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL	
	Midline	3 mL	3 mL		3 mL	
	Implanted Port	5 mL	10 mL	10 mL		5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL	
	Tunneled & non-Tunneled Catheters.	5 mL	10 mL	10 mL	5 mL	
<b>PEDIATRIC 33 LBS - 66LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL	
	Midline	3 mL	3 mL		3 mL	
	Implanted Port	5 mL	5 mL	10 mL	3 mL	
	Peripherally Inserted Central Catheter (PICC)	5 mL	5 mL	10 mL	3 mL	
	Tunneled & non-Tunneled Catheters.	5 mL	5 mL	10 mL	3 mL	

**FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

*\*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.*