

IRON DEFICIENCY ANEMIA REFERRAL FORM

PATIENT DEMOGRAPHICS:

PATIENT NAME:	PATIENT'S CONTACT #:
DATE OF REFERRAL:	ADDRESS:
DATE OF BIRTH:	CITY, STATE, ZIP:
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:	
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE MALE
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST NKDA

PRIMARY DIAGNOSIS:

D50.0 Iron Deficiency Anemia	D50.8 Other Iron Deficiency Anemias
D50.9 Iron Deficiency Anemia, Unspecified	D63.1 Anemia in chronic kidney disease
N18.1 Chronic kidney disease, stage 1	N18.2 Chronic kidney disease, stage 2
N18.3 Chronic kidney disease, stage 3	N18.4 Chronic kidney disease, stage 4

REQUIRED DOCUMENTATION: Please provide a copy of the following documents.

<input checked="" type="checkbox"/> 1. INSURANCE CARD (Front & Back)	<input checked="" type="checkbox"/> 2. PATIENT DEMOGRAPHICS	<input checked="" type="checkbox"/> 3. MOST RECENT LABS	<input checked="" type="checkbox"/> 4. MEDICATION LIST
<input checked="" type="checkbox"/> 5. H & P	<input checked="" type="checkbox"/> 6. BONE SCAN	<input checked="" type="checkbox"/> 7. MOST RECENT FERRITIN, HEMOGLOBIN, AND HEMATOCRIT	<input checked="" type="checkbox"/> 8. OTHER IRON STUDIES
<input checked="" type="checkbox"/> Intolerance or unsatisfactory response to oral? (Provide documentation)		Y	N

PRIMARY MEDICATION ORDER:

PRN & PREMEDICATIONS:

Injectafer 750 mg IV on day 1 and day 7 (for patients 50 kg or greater). Injectafer 15 mg/kg IV (not to exceed 1,500 mg) on day 1 and day 7 (for any weight). Monoferic 20 mg/kg IV once (for patients weighing less than 50 kg). Monoferic 1,000 mg IV once (for patients weighing greater than 50 kg). Venofer 200 mg IV every 2-3 days for 5 doses. Venofer 300 mg IV every 3 days for 3 doses. Feraheme 510 mg IV on day 1 and day 3. INFeD _____ mg IV over 4 hours. Other: _____ FIRST DOSE: Y N <input checked="" type="checkbox"/> Based on payer guidelines, Venofer may be substituted for Injectafer. <input checked="" type="checkbox"/> Refill x12 months unless otherwise noted.	MEDICATIONS	30 minutes prior to every infusion	PRN
	Acetaminophen 650 mg PO		PRN every ___ hour for mild or moderate infusion reaction.
	Diphenhydramine 25 mg PO		PRN every ___ hour for mild or moderate infusion reaction.
	Diphenhydramine 25 mg IV		PRN every ___ hour for mild or moderate infusion reaction.
	Methylprednisolone 125 mg IV		PRN every ___ hour for mild or moderate infusion reaction.
	Other: _____		PRN every ___ hour for mild or moderate infusion reaction.

LINE USE/CARE ORDERS:

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

<input checked="" type="checkbox"/> START PIV/ACCESS CVC <input checked="" type="checkbox"/> FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE) OTHER FLUSH ORDERS: (please fax other reaction orders if checking this box)	<input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) OTHER: (please fax other reaction orders if checking this box)
---	---

PRESCRIBER INFORMATION: Please check preferred form of communication.

PHYSICIAN NAME:	PHONE:
OFFICE CONTACT:	FAX:
ADDRESS:	EMAIL:
CITY, STATE, ZIP:	NPI:

➔ (GENERIC SUBSTITUTION PERMITTED)

PHYSICIAN SIGNATURE:

DATE:

➔ (DISPENSE AS WRITTEN)

PHYSICIAN SIGNATURE:

DATE:

IRON DEFICIENCY ANEMIA REFERRAL FORM

FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

**This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
SYMPTOM CLASSIFICATION	<ul style="list-style-type: none"> Flushing Dizziness Headache Apprehension Diaphoresis Palpitations Nausea / Vomiting Pruitis 	<ul style="list-style-type: none"> Chest Tightness Shortness of Breath Hypo/hypertension (>20 mmHg Change in Systolic BP from Baseline) Increased Temperature (>2 Degrees Fahrenheit) Urticaria 	<ul style="list-style-type: none"> Hypo/hypertension (>40 mmHg Change in Systolic BP from Baseline). Increase Temperature (>2 Degrees Fahrenheit) with Rigors Shortness of Breath with Wheezing Laryngeal Edema Chest Pain Hypoxemia
TREATMENT PROTOCOL FOR ADULTS >66LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
ADULT > 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
PEDIATRIC 33 LBS - 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.