



## OCREVUS REFERRAL FORM

| PATIENT DEMOGRAPHICS:   |   |   |   |  |
|---|---|---|---|--|
| PATIENT NAME:   | PATIENT'S CONTACT #:  |   |   |  |
| DATE OF REFERRAL:   | ADDRESS:  |   |   |  |
| DATE OF BIRTH:  | CITY, STATE, ZIP:   |   |   |  |
| FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:   |   |   |   |  |
| HEIGHT: _____ FEET      _____ INCHES  | GENDER:      FEMALE                      MALE   |   |   |  |
| WEIGHT: _____ LB      or      _____ KG  | ALLERGIES:      SEE LIST                      NKDA  |   |   |  |
| PRIMARY DIAGNOSIS:  |   |   |   |  |
| ICD-10 CODE:      G35 - Multiple sclerosis                      Other   |   |   |   |  |
|   |   |   |   |  |
|   |   |   |   |  |
| REQUIRED DOCUMENTATION: Please provide a copy of the following documents.   |   |   |   |  |
| <input checked="" type="checkbox"/> 1. INSURANCE CARD (Front & Back) <input checked="" type="checkbox"/> 2. PATIENT DEMOGRAPHICS <input checked="" type="checkbox"/> 3. MOST RECENT LABS <input checked="" type="checkbox"/> 4. MEDICATION LIST<br><input checked="" type="checkbox"/> 5. H & P <input checked="" type="checkbox"/> 6. MRI RESULTS <input checked="" type="checkbox"/> 7. NEGATIVE HEPATITIS B, SEROLOGY <input checked="" type="checkbox"/> 8. IMMUNOGLOBULINS PANEL |   |   |   |  |
| PRIMARY MEDICATION ORDER:   |   |   |   |  |
| Please include MEDICATION, DOSE, FREQUENCY, DURATION and any ADDITIONAL administration INSTRUCTIONS specific to the primary therapy.<br><br>Loading Dose: Ocrevus 300 mg IV on day 1 and day 15.<br>Maintenance Dose: Ocrevus 600 IV mg every 6 months.<br>Other:<br><br>FIRST DOSE:      Y                      N<br><input checked="" type="checkbox"/> Refill x12 months unless otherwise noted.   | <th colspan="3">PRN &amp; PREMEDICATIONS:</th>  | PRN & PREMEDICATIONS:                     |   |  |
|   | <b>MEDICATIONS</b>  | <b>30 minutes prior to every infusion</b> | <b>PRN</b>  |  |
|   | Acetaminophen 650 mg PO   |   | PRN every ____ hour for mild or moderate infusion reaction. |  |
|   | Diphenhydramine 25 mg PO  |   | PRN every ____ hour for mild or moderate infusion reaction. |  |
|   | Diphenhydramine 25 mg IV  |   | PRN every ____ hour for mild or moderate infusion reaction. |  |
|   | Methylprednisolone 125 mg IV  |   | PRN every ____ hour for mild or moderate infusion reaction. |  |
|   | Other: _____  |   | PRN every ____ hour for mild or moderate infusion reaction. |  |
| LINE USE/CARE ORDERS:   |   |   |   |  |
| <input checked="" type="checkbox"/> START PIV/ACCESS CVC<br><input checked="" type="checkbox"/> FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE)<br>OTHER FLUSH ORDERS: (please fax other reaction orders if checking this box)   | <th colspan="3">ADVERSE REACTION &amp; ANAPHYLAXIS ORDERS:</th>   | ADVERSE REACTION & ANAPHYLAXIS ORDERS:    |   |  |
|   | <input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side)<br>OTHER: (please fax other reaction orders if checking this box) |   |   |  |
| PRESCRIBER INFORMATION: Please check preferred form of communication.   |   |   |   |  |
| PHYSICIAN NAME:   | PHONE:  |   |   |  |
| OFFICE CONTACT:   | FAX:  |   |   |  |
| ADDRESS:  | EMAIL:  |   |   |  |
| CITY, STATE, ZIP:   | NPI:  |   |   |  |
|  (GENERIC SUBSTITUTION PERMITTED)<br>PHYSICIAN SIGNATURE: _____  |   | DATE: _____                               |   |  |
|  (DISPENSE AS WRITTEN)<br>PHYSICIAN SIGNATURE: _____   |   | DATE: _____                               |   |  |

# OCREVUS REFERRAL FORM

## FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

*\*This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

|   | MILD INFUSION REACTION   | MODERATE INFUSION REACTION   | SEVERE INFUSION REACTION/ANAPHYLAXIS   |
|---|--|--|--|
| <b>SYMPTOM CLASSIFICATION</b>                         | <ul style="list-style-type: none"> <li>Flushing</li> <li>Dizziness</li> <li>Headache</li> <li>Apprehension</li> <li>Diaphoresis</li> <li>Palpitations</li> <li>Nausea / Vomiting</li> <li>Pruitis</li> </ul> | <ul style="list-style-type: none"> <li>Chest Tightness</li> <li>Shortness of Breath</li> <li>Hypo/hypertension (&gt;20 mmHg Change in Systolic BP from Baseline)</li> <li>Increased Temperature (&gt;2 Degrees Fahrenheit)</li> <li>Urticaria</li> </ul> | <ul style="list-style-type: none"> <li>Hypo/hypertension (&gt;40 mmHg Change in Systolic BP from Baseline).</li> <li>Increase Temperature (&gt;2 Degrees Fahrenheit) with Rigors</li> <li>Shortness of Breath with Wheezing</li> <li>Laryngeal Edema</li> <li>Chest Pain</li> <li>Hypoxemia</li> </ul>   |
| <b>TREATMENT PROTOCOL FOR ADULTS &gt;66LBS</b>        | <input checked="" type="checkbox"/> Administer PRN medications per Physician order   | <input checked="" type="checkbox"/> Administer PRN medications per Physician order   | <input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting.<br><input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access.<br><input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed.<br><input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.   |
| <b>TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS</b> | <input checked="" type="checkbox"/> Administer PRN medications per Physician order   | <input checked="" type="checkbox"/> Administer PRN medications per Physician order   | <input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting.<br><input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access.<br><input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min<br><input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed.<br><input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1. |

**FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

### FLUSHING PROTOCOLS

|                                      |  | FLUSHING PROTOCOL<br>Normal Saline* |            | LOCKING PROTOCOL<br>Heparin Sodium |                |
|--------------------------------------|--|-------------------------------------|------------|------------------------------------|----------------|
|                                      |  | 0.9% Sodium Chloride                |            | 10 Units/mL                        | 100 Units/mL   |
| PATIENT CLASSIFICATION               | LINE TYPE                                      | PRE-ADMIN                           | POST ADMIN | POST LAB DRAW                      | POST NS FLUSH* |
| <b>ADULT &gt; 66 LBS</b>             | Peripheral IV Catheter                         | 3 mL                                | 3 mL       |                                    | 3 mL           |
|                                      | Midline  | 3 mL                                | 3 mL       |                                    | 3 mL           |
|                                      | Implanted Port                                 | 5 mL                                | 10 mL      | 10 mL                              | 5 mL           |
|                                      | Peripherally Inserted Central Catheters (PICC) | 5 mL                                | 10 mL      | 10 mL                              | 5 mL           |
|                                      | Tunneled & non-Tunneled Catheters              | 5 mL                                | 10 mL      | 10 mL                              | 5 mL           |
| <b>PEDIATRIC<br/>33 LBS - 66 LBS</b> | Peripheral IV Catheter                         | 3 mL                                | 3 mL       |                                    | 3 mL           |
|                                      | Midline  | 3 mL                                | 3 mL       |                                    | 3 mL           |
|                                      | Implanted Port                                 | 5 mL                                | 5 mL       | 10 mL                              | 3 mL           |
|                                      | Peripherally Inserted Central Catheters (PICC) | 5 mL                                | 5 mL       | 10 mL                              | 3 mL           |
|                                      | Tunneled & non-Tunneled Catheters              | 5 mL                                | 5 mL       | 10 mL                              | 3 mL           |

**FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

\*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.